

Persistent spin and mass currents and Aharonov-Casher effect

A. V. Balatsky

*Center for Materials Science and Theoretical Division, T-11,
Los Alamos National Laboratory, Los Alamos, NM 87545 and
Landau Institute for Theoretical Physics, Moscow, Russia*

B. L. Altshuler

Department of Physics, MIT, Cambridge, MA 02139

(Received February 1, 2008)

Spin-orbit interaction produces persistent spin and mass currents in the ring via the Aharonov-Casher effect. The experiment in $^3\text{He} - A_1$ phase, in which this effect leads to the excitation of mass and spin supercurrent is proposed.

PACS Nos. 71.70.Ej; 03.65.Bz; 67.50.Fi.

The discovery in 1983 of the Berry phase [1] brings a new understanding of different topological effects in quantum mechanics. The simplest realization of Berry phase is Aharonov-Bohm (AB) effect of charged particle in external electromagnetic field. A transparent demonstration of the AB effect is a persistent current in mesoscopic rings threaded by magnetic field [2], as well as many other experiments, which proved the relevance of AB effect on transport in multiconnected geometry. More recently it has been pointed out that there is an analog of AB effect in the presence of the spin-orbit (SO) interactions, which leads to nontrivial phase shifts and to topological interference effect of the wave function of a particle with spin, which was called Aharonov-Casher (AC) effect [3].

There is a major difference between AB and AC effects: the AB effect comes from the true gauge invariant coupling $j_\mu A_\mu$ between the current j_μ and the electromagnetic vector potential A_μ and for that reason can be observed even in the absence of magnetic field $\mathbf{B} = \text{curl}\mathbf{A}$ in the region where particles are propagating. In the AC effect, on the other hand, the phase of the wave function is changed as a result of SO interaction i.e. as a coupling of the spin current $j_\mu^{\sigma_i}$ to an effective tensor gauge potential $E_\nu \epsilon_{\mu\nu i}$ where $\epsilon_{\mu\nu\lambda}$ is the antisymmetric tensor, and $\mathbf{E}$ is the electric field. Therefore nonzero electric field on the way of the particle is necessary in order to produce the AC phase shift [4]. In the case of electrons moving in the atomic electric field SO coupling can be written in a more familiar form as $\boldsymbol{\sigma}\mathbf{l}$ with $\mathbf{l}$ being the orbital momentum of electron [5].

Inspite this important difference the unifying point of view on both effects is that they are consequence of the Berry phase acquired by the wave function of the particle under adiabatic transport of the particle from some initial state through the set of intermediate states back to its original configuration.

Here we will implement this point of view and report on the observation that AC effect can result in persistent spin and mass currents. The wave function of a particle in external magnetic and electric fields and in the presence of SO interaction will acquire the spin dependent Berry phase: i) additional phase of the wave function is given by $\phi = \Phi_{AB} + \sigma_z \Phi_{AC}$, where Φ_{AB} is the AB flux piercing the ring (in case if particles are charged) and

Φ_{AC} is the AC flux due to AC effect, $\sigma_z = \pm$ is the spin projection (we assume spin $\frac{1}{2}$ particles). The formerly doubly degenerate eigenstates and eigenvalues will acquire spin dependent shift $E_n(\Phi_{AB} + \sigma_z \Phi_{AC})$, $\Psi_n(\Phi_{AB} + \sigma_z \Phi_{AC})$, see also [6]; ii) this spin dependent shift will lead to a persistent spin current:

$$j_\varphi^{\sigma_z} = -\frac{c}{4\pi R} Tr \frac{\partial E(\Phi_{AB} + \sigma_z \Phi_{AC})}{\partial \Phi_{AC}} \sigma_z = -\frac{c}{2\pi R} \frac{\partial E}{\partial \Phi_{AB}} \quad (0.1)$$

with $j_\varphi^{\sigma_z}$ being the z component of the spin current along the ring, σ_i are the Pauli matrices. In the presence of the net spin polarization AC effect also leads to mass current proportional to $n_\uparrow - n_\downarrow$.

Another realization of the Berry phase leading to persistent spin currents was discussed in [7]. Note that the state with persistent spin current does not violates P (parity) and T (time reversal) invariance, in the contrast to persistent mass current. Due to this reason persistent spin current can be excited in the absence of external magnetic flux. Actually for an electron state with nonzero angular momentum ℓ (say in an atom) persistent spin current means nothing but existence of the SO interaction; iii) the AC effect and persistent spin currents are independent on the charge of the particle and can be observed for neutral particles , as was done in original observation of the AC effect for neutrons [8]; iv) the time dependent AC flux generates effective spin dependent “electric” field $\mathbf{E}_{AC}$ via the Faraday law $\frac{1}{c} \partial_t \Phi_{AC} = - \oint \mathbf{E}_{AC} d\mathbf{l}$ acting on neutral as well as on charged spinfull particles. This is the local effect,

independent on the macroscopic phase coherence [9].

The closely related, however different, problem on the effect of random SO interactions on the electron transport properties in mesoscopic ring was considered by Meir, Gefen and Entin-Wohlman [6]. They showed that in the presence of SO scattering the flux dependence of energy and eigenfunctions acquires the spin dependent shift $\Phi_{AB} \pm \delta$ where shift δ is governed by an averaged over the ring random SO interactions. Thus the overall effect of SO interaction in this approach is given by the particular impurity configuration and independent of external fields. In our case we will neglect the random SO scattering and

consider the AC effect in external electric field, which can be varied in the direction and magnitude. This will lead to spin current excitation due to Faraday law for time dependent AC flux. Interesting realization of this effect will be shown below to take place in superfluid ^3He , where external crossed electric and magnetic field should cause a supercurrent.

Consider first a one dimensional ring of radius R . We will describe it using a tight binding model on a closed chain of N sites separated by a distance $a = R/N$. An external magnetic field perpendicular to the plane $\mathbf{B} \parallel \mathbf{e}_z$ (we will use cylindrical coordinates given by unit vectors $\mathbf{e}_\phi, \mathbf{e}_\rho, \mathbf{e}_z$) results in the twisted boundary conditions for a charged particle wave function $\Psi(N) = \exp(i2\pi\Phi_{AB}/\Phi_0)\Psi(0)$ with the AB flux $\Phi_{AB} = B_z\pi R^2$ and $\Phi_0 = \frac{hc}{e}$. The Hamiltonian of this chain taking into account the SO interaction and Zeeman splitting $g\mu_B\hat{\boldsymbol{\sigma}}\mathbf{B}$ (g being the gyromagnetic ratio) can be written as

$$H = -t \sum_{n,\sigma,\sigma'} \hat{\Lambda}(n) |n, \sigma\rangle \langle n+1, \sigma'| + h.c. + \sum_{n,\sigma,\sigma'} (\epsilon_n + g\mu_B\hat{\boldsymbol{\sigma}}\mathbf{B}) |n, \sigma\rangle \langle n, \sigma'| \quad (0.2)$$

here n labels sites, ϵ_n denotes the onsite energies, and $\hat{\Lambda}_{\sigma\sigma'}(n) = \exp(i\hat{A}_{AC}(n, n+1))_{\sigma\sigma'} = \exp(i\frac{q\mu_B}{hc}\hat{\boldsymbol{\sigma}} \int_n^{n+1} d\mathbf{r} \times \mathbf{E}(\mathbf{r}))_{\sigma\sigma'}$, where $\hat{A}_{AC}$ is the AC analog of the vector potential $\Phi_{AC} = \sum_{n=1}^{N-1} A_{AC}(n, n+1)$.

Following the approach of Ref [6] we use the transfer matrix T_N defined as:

$$\begin{pmatrix} \psi_N \\ \psi_{N-1} \end{pmatrix} = T_N \begin{pmatrix} \psi_1 \\ \psi_0 \end{pmatrix} \quad (0.3)$$

where ψ_n is the spinor wave function of the particle in coordinate representation, and we drop out the spin indexes for simplicity. Using the structure of the Hamiltonian Eq(2) we can write T_N as a direct product:

$$T_N = \hat{S} \otimes T'_N, T'_N = t^N \begin{pmatrix} 1 & 0 \\ \sum_{i=1}^N (\epsilon_i + g\mu_B\sigma_z) & 1 \end{pmatrix} \quad (0.4)$$

$$\hat{S} = \prod_{i=1}^N \hat{\Lambda}(i) \quad (0.5)$$

To derive this equation we had to neglect the noncommutativity of $[\hat{\Lambda}, g\mu_B B\sigma_z] \sim O(B \cdot E)$, which leads to higher order in the external fields contributions. As we will see the Berry phase will be proportional to the circulation of E_ρ on the ring; thus taking into account commutator will lead to higher order terms. Strictly speaking Eqs(4,5) are valid in the $N \rightarrow \infty$ limit when $\hat{\Lambda}(0) = \hat{\Lambda}(N)$.

For the slowly varying electric field $\mathbf{E}$ ($|\partial_\phi \mathbf{E}|/|\mathbf{E}| a \ll 1$, where a is a distance between sites) the spin transfer matrix $\hat{S}$ equals to:

$$\begin{aligned} \hat{S} &= \prod_{i=1}^N \exp(i \frac{g\mu_B}{\hbar c} \hat{\sigma} \int_i^{i+1} d\mathbf{r} \times \mathbf{E}(\mathbf{r})) \\ &\cong \exp(i \frac{g\mu_B}{\hbar c} \hat{\sigma} \oint d\mathbf{r} \times \mathbf{E}(\mathbf{r})) \end{aligned} \quad (0.6)$$

We neglected the commutator in the exponent in Eq (6) coming from the Hausdorff's formula $e^A e^B = e^{A+B+1/2[A,B]}$, since it is proportional to $\partial_\phi \mathbf{E}$. Only radial component of electric field E_ρ contributes to the contour integral in Eq (6). As a result

$$\frac{g\mu_B}{\hbar c} \oint d\mathbf{r} \times \mathbf{E}(\mathbf{r}) = -\mathbf{e}_z 2\pi \frac{\Phi_{AC}}{\Phi_0} \quad (0.7)$$

using Eqs (6,7) the eigenfunctions equation with twisted boundary conditions can be written in the compact form

$$T'_N \begin{pmatrix} \psi_1 \\ \psi_0 \end{pmatrix} = \exp(2\pi i \frac{\Phi_{AB}}{\Phi_0} + 2\pi i \sigma_z \frac{\Phi_{AC}}{\Phi_0}) \begin{pmatrix} \psi_1 \\ \psi_0 \end{pmatrix} \quad (0.8)$$

It is obvious that in this geometry spin dependent AC flux coming from SO interactions enters the transfer matrix as a spin dependent phase. This equation allows to find energy eigenvalues and eigenfunctions of the tight binding Hamiltonian

Eq (2) if they are known for the bare problem without SO interactions. Namely, the energy spectrum and the wave functions will depend on the effective flux which is a sum of spin-independent (AB) and spin dependent (AC) parts: $E_n = E_n(\Phi_{AB} + \sigma_z \Phi_{AC})$, $\Psi_n = \Psi_n(\Phi_{AB} + \sigma_z \Phi_{AC})$, as was mentioned. The energy dependence on the Φ_{AC} leads to the persistent spin current $j_\phi^{\sigma_z}$ (see Eq (1)). Note again that the spin dependent Berry phase

Φ_{AC} is nonzero even for neutral particles. This equivalence of the AC and AB effect holds only if the spin relaxation is neglected.

It is instructive to estimate the magnitude of the AC flux for realistic mesoscopic systems. From Eq (7) it follows:

$$\frac{\Phi_{AC}}{\Phi_0} = g\mu_B \frac{RE}{c\hbar} \quad (0.9)$$

The relativistic nature of the AC effect is reflected in the ratio of electric potential on the scale of the size of the ring eRE to the rest energy of the particle mc^2 .

If the electric field is time dependent the variation of the AC flux results in the appearance of a spin-dependent driving force due to Faraday law. This causes the spin current in accord with Ohm's law. Applying Faraday law to spin $\uparrow\downarrow$ liquids (assuming $\Phi_{AB} = const$) we find that particles experience spin dependent force $-\sigma_z \frac{1}{c} \partial_t \Phi_{AC} = \sigma_z \oint \mathbf{E}_{AC} d\mathbf{l}$. Again, in the presence of net spin polarization this force will cause mass current. This effect is a consequence of the Lorentz invariance and electrodynamics and is a local phenomenon, as it has been pointed out previously [9]. In order to excite spin and mass current due to local force the global phase coherence along the ring *is not required*.

For the ring of the radius $R \sim 10^{-3}$ cm, in the external electric field $E \sim 10^5$ v/cm Eq(9) gives for particles with gyromagnetic ratio $g \sim 1$ that $\frac{\Phi_{AC}}{\Phi_0} \sim 10^{-3}$ (see Eq(9)), that is a tiny effect. On the other hand, in the semiconductors effective g -factor can be two orders of magnitude larger [10]. For these samples effective flux will be of the order of $10^{-1}\Phi_0$, which makes the interference effects associated with AC effect in external fields experimentally observable. Still the experimental observation of the effect in a real mesoscopic ring does not look as a simple problem. The main difficulties are due to the screening of the electric field and the necessity to work with considerably strong magnetic fields which prevent the usage of a SQUID.

Consider another class of systems with phase coherence established on the macroscopic scale – superfluid 3He . In strong enough magnetic field ($B \sim 10kG$) superfluid transition is known to split on two phase transitions [11]: 1) First atoms with spins along the field are

paired – this is called ${}^3\text{He} - A_1$ phase; 2) After second phase transition atoms of both spin directions are paired in ${}^3\text{He} - A_2$ phase.

${}^3\text{He} - A_1$ phase is a superfluid with $S_z = +1$ Cooper pairs condensate. Atoms with spins opposite to the magnetic field are not condensed. In certain range of temperatures $\Delta T \sim 6 \cdot 10^{-3} \text{mK}/\text{kG} \cdot B$ the ${}^3\text{He} - A_1$ is the only superfluid phase. Thus the ${}^3\text{He} - A_1$ phase has important features: 1) the phase coherence of the of $S_z = +1$ condensate over the macroscopic distances and 2) total spin polarization of the coherent subsystem. This implies that in external electric field condensate will exhibit AC effect.

We propose the following experiment. Consider ${}^3\text{He} - A_1$ phase within a capacitor with the plates lying in the (xy) plane, so that electric field has only z component. Let the spins be polarized along y axis. In this case the phase of the condensate wave function acquires the position dependent shift $\phi(x) = \phi(0) + 2\pi \frac{\Phi_{AC}(x)}{\Phi_0}$, with $\Phi_{AC}(x)$ given by Eq(7) with integral taken along x axis. This generates the condensate flow with the velocity $v_s = \frac{\hbar}{2m_3} \partial_x \phi(x)$, i.e. supercurrent. For $E \sim 10^5 \text{V}/\text{cm}$ this current will be $v_s \simeq 2 \cdot 10^{-7} \text{cm}/\text{sec}$ (compare with the critical velocity in ${}^3\text{He} - A$ $v_c \simeq 0.02 \text{cm}/\text{sec}$ [11]). The phase difference accumulated by the Cooper pair upon transport through the capacitor of length L will be $\Delta\phi \simeq 6 \cdot 10^{-4} L[\text{cm}]$. In order to have a phase difference $\Delta\phi \simeq 0.1 \cdot 2\pi$ we need to have a channel! of length $L \sim 1\text{m}$. Presently in

most experiments on ${}^3\text{He}$ are done in containers with characteristic size of few cm .

The supercurrent flow can be detected experimentally since it causes the normal counterflow due to mass conservation. The latter leads to the heat flow $\dot{Q}$ opposite to the supercurrent [12]:

$$\dot{Q} = (v_s - v_n)(\rho_s/\rho)(TS)(Area) \quad (0.10)$$

where v_s and v_n are superfluid and normal velocities, ρ_s and ρ stand for superfluid and total densities and S is the ${}^3\text{He}$ entropy per unit volume. Using Eq (10) we can estimate $\dot{Q}$ at $E \sim 10^5 \text{V}/\text{cm}$ ($v_s \sim 10^{-7} \text{cm}/\text{sec}$) as $\dot{Q}/\text{area} \simeq 10^{-8} \text{erg}/\text{sec}$.

The ac modification of the same idea looks even more realistically. We propose the

experiment in the capacitor with electric field oscillating with some frequency ω . Such a field will cause oscillating super and therefore normal currents which apparently can be detected with less difficulties. The power loss is known [13] to be equal to $P \simeq 8\pi L\eta v_s^2$, where L is the length of the channel and $\eta \sim 3 \cdot 10^{-2} \frac{g}{cmsec}$ is the normal He^3 viscosity. The estimation for the field $E \sim 10^5 V/cm$ gives $P \simeq 10^{-14} erg/sec$. For example, the frequency dependence of the impedance of the capacitor can be measured with the high accuracy. Due to viscous flow of the normal component the impedance in ${}^3He - A_1$ will be different from the both normal and any other superfluid phase : only in A_1 phase the AC effect leads to the mass supercurrent and therefore to the normal component flow.

To conclude , we proposed the new realization of the AC effect, which leads to persistent spin and mass currents in mesoscopic ring and in the superfluid ${}^3He - A_1$ phase. We argue that time dependent electric field in particular geometry will excite *locally* spin current even in the macroscopic samples via the Faraday law.

We acknowledge useful discussions with A. Leggett, D. Osheroff and G. Swift. This work was supported by J. R. Oppenheimer fellowship and by Department of Energy (A.B.), by Joint Services Electronic Program Contract No DAAL03-89-0001 (B.A.) and by Advanced Studies Program of the Center for Materials Science at LANL.

REFERENCES

- [1] M. V. Berry, Proc. of Royal Soc. London, **A 392**, 45, (1984).
- [2] M. Büttiker, Y. Imry and R. Landauer, Phys. Lett., **96 A**, 365, (1983); L. P. Levy, G. Dolan, J. Dunsmuir and H. Bouchiat, Phys. Rev. Lett., **64**, 2094, (1990).
- [3] Y. Aharonov and A. Casher, Phys. Rev.Lett.,**53**, 319, (1984).
- [4] A. Goldhaber, Phys. Rev. Lett., **62**, 482, (1989).
- [5] V. B. Berestetskii, E. M. Lifshitz and L. P. Pitaevskii, " Quantum Electrodynamics", (Pergamon Press, 1982), §42.
- [6] Y. Meir, Y. Gefen and O. Entin-Wohlman, Phys. Rev. Lett., **63**, 798, (1989).
- [7] D. Loss, P. Goldbart and A. V. Balatsky, Phys. Rev. Lett, **65**, 1655, (1990).
- [8] A. Cimmino et. al., Phys. Rev. Lett., **63**, 380, (1989).
- [9] The analogous effect for Berry phase due to Zeeman coupling was pointed out by A. Stern, Phys. Rev. Lett., **68**, 1022, (1992).
- [10] H. Mathur and A. D. Stone, Phys. Rev. Lett., **68**, 2964, (1992).
- [11] A. J. Leggett, Rev. Mod. Phys., **47**, 331, (1975). See Sec. 7, 10.
- [12] J. Wheatley, Rev. Mod. Phys., **47**, 436, (1975).
- [13] G.K. Batchelor, "An Introduction to Fluid Dynamics", Cambridge Univ. Press, 1977.